# Who should get screened?

- » Anyone over 45 years or if strong family history of CRC or certain types of polyps.
- » History of intestinal polyps.
- » Anyone who suffers from IBS.
- » Personal history of breast cancer.

# You can prevent.

There is no reason to wait for symptoms to appear, waiting may be too late.

### **Risk factors:**

- » Being over 45 years old.
- » High consumption of red or processed meats.
- » Diet low in fiber, fruits and vegetables.
- » Sedentary.
- » Overweight and obesity.
- » Consumption of alcohol.
- » Smoking.
- » Intestinal polyps.
- » Inflammatory bowel disease.









Preserving the health and quality of life of people.

Contact us for more information about the

OC-Auto FIT-CHEK™ Screening Method for Hemoglobin in Stool.

#### **CDMX & Metropolitan area**

Office: +52 (55) 5294 7193 WhatsApp: +52 1 (55) 7381 9171 info@MedPrimex.com

Get to know our products:

www.MedPrimex.com








The test is manufactured in U.S.A. in an ISO 13485 quality system in a facility registered by the FDA to guarantee reliability and reproducibility.



OC-Auto FIT-CHEK

Colorectal cancer is treatable as long as it is detected early.

Prevention is the best strategy for staying healthy.









# Colorectal cancer is treatable as long as it is detected early.

OC-Auto FIT-CHEK™ is a NON-INVASIVE method that does not require prior preparation, special diet, intestinal lavage, nor hospitalization. The sample can be collected at any time from the comfort of your home

90% OF THE CASES HAVE CURE, IF IT IS DETECTED EARLY.

# Signs and symptoms:

- » Usually appear in advanced stages.
- » Changes in bowel habits (diarrhea, constipation, thinness)
- » Bleeding, dark stools, blood in stool.
- » Abdominal colic
- » Need to defecate that does not disappear after doing so..

Technology based on the immunological screening method for hemoglobin in stool, that allows for a timely, rapid, and economic detection of CRC, before symptoms appear, decreasing the mortality rate and avoiding false negatives or false positive results.







**Anatomy**Lower digestive system

# Important facts



It is the 3<sup>rd</sup> most common cancer in the world & the 4<sup>th</sup> most common cause of death.

It is the 4<sup>th</sup> most frequent type of cancer in Mexico.



In Mexico it represents the 2<sup>nd</sup> cause of death associated with cancer. It originates in the rectum or colon, with the growth of a non-cancerous tumor.



This precancerous phase is totally asymptomatic, but detectable.

90 out of every 100 people survive after being diagnosed in the early stages.



70% of those affected go to consultation when the disease is irreversible